



PRIMHD summary report – HoNOS65+

Health of the Nation Outcomes Scales – adults 65+ report for New Zealand



tepou.co.nz

This report summarises national Programme for the Integration of Mental Health Data (PRIMHD) data submitted by Te Whatu Ora Districts. In particular, it presents Health of the National Outcomes Scale for older adult (HoNOS65+) data from services where HoNOS65+ is the primary measure.

This report is organised into three major sections that provide information about:

- 1. Outcomes related information:** This provides indications about what changes have occurred for service users between entering and leaving the service. Outcome is assessed by comparing the group admitted and the group discharged from the service in the same time period. This should provide a reasonable indication of outcomes achieved unless the service user mix has changed significantly over the usual period for which service is delivered.
- 2. Service related information:** This provides information about the services, such as the overall severity of service users who use different services.
- 3. Collection completion and validity:** This details the completeness of the data set provided by your mental health services. This is important because it indicates how valid and reliable the data sets are likely to be. The less complete the data set, the less valid the information is likely to be.

In many cases the data is presented graphically for New Zealand, and then presented as a table for the individual team types (see team type classification factsheet).

The time period covered differs for the different data presented. See the title or the notes under the graphs or tables for information about the time period covered. Unless otherwise stated, the notes under the graphs also apply to the corresponding table.

For all graphs and tables, if there are less than twenty cases in the data set, then the information is not presented. This is because small samples frequently provide inaccurate and potentially misleading results. See the notes and user guide for other important information about the graphs and tables.

Where appropriate, the statistical **confidence interval** is presented. This is shown by error bars (small lines above and below the average) on the graphs, and a score range in some tables. As a rule of thumb, if the confidence intervals of two data points do not overlap, the two points can be considered to be significantly different. If the confidence intervals of the data points do overlap, we assume the points are not significantly different. It is important to note that statistical significance may not indicate a clinically significant difference.

See the associated user guide for more information about how to understand and use the data presented in this report.


Data for tables and graphs was extracted 26 January 2026 from PRIMHD by Te Whatu Ora, then analysed and formatted by Te Pou.

Please note: For this period a few Te Whatu Ora Districts have incomplete data which will affect New Zealand totals.

1. Outcomes – changes in service user status

This section presents HoNOS65+ data indicating the status of service users at different stages of their contact with Te Whatu Ora District mental health services. Graph and Table 1, and Graph 2 show results from HoNOS65+ total scores. Graphs and Tables 3 to 5b show results related to the percentage of HoNOS65+ items in the clinical range. Table 6 shows results from the Index of Severity derived from HoNOS65+ scores.

Graph 1: Average HoNOS65+ total score (12 items) by collection type: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025



Notes: Error bars indicate the confidence intervals around the data point. If error bars overlap, the data points are not significantly different. Community discharge does not include discharge to an inpatient unit.

Interpretation: Decrease between admission and discharge is an indication of the outcomes achieved by the service user and service. The greater the decrease between admission and discharge, and the lower the average HoNOS65+ score at discharge, the more positive the outcome.

Target: A greater decrease from admission to discharge and lower average rating at discharge.


Table 1: Average HoNOS65+ total score (12 items), by collection type and team: New Zealand, Oct 2025 - Sep 2025

Team type	Assessment only			Admission			Review			Discharge		
	N	Mean	CI	N	Mean	CI	N	Mean	CI	N	Mean	CI
Community services												
Alcohol and drug team							39	6.6	5.2-8.0			
Community team	349	11.2	10.6-11.8	201	12.3	11.5-13.2	1,708	9.2	9.0-9.5	166	7.8	6.9-8.6
Kaupapa Māori team							56	15.3	13.5-17.1			
Needs assessment and service coordination team							44	8.3	7.0-9.7			
Older people team	959	11.0	10.6-11.4	2,253	10.6	10.3-10.8	3,325	7.8	7.6-8.0	1,634	7.1	6.8-7.3
Pacific people team							26	5.5	3.9-7.1			
Specialty team	220	13.2	12.4-14.0	337	16.0	15.3-16.6	503	13.6	13.1-14.0	343	12.0	11.5-12.5
Total	1,541	11.3	11.1-11.6	2,816	11.4	11.1-11.6	5,714	8.8	8.7-8.9	2,160	7.9	7.7-8.1
Inpatient services												
Forensic team							25	8.5	6.0-11.0			
Inpatient team				336	14.4	13.6-15.1	44	10.4	8.1-12.7	219	8.9	8.0-9.8
Older people team				739	15.8	15.3-16.2	144	12.7	11.6-13.8	586	8.1	7.6-8.6
Total				1,084	15.3	14.9-15.7	213	11.7	10.8-12.7	814	8.3	7.9-8.7

Notes: N = number of collections in period. Average = average HoNOS65+ (12 item) score, CI = confidence interval for average score. Community discharge does not include discharge to an inpatient unit.

Interpretation: If confidence intervals for two scores do not overlap, then the scores have a statistically significant difference. It is important to note that statistical significance may not indicate a clinically significant difference.

Graph 2: Difference in HoNOS65+ total score (12 items) for matched pairs by pair type and setting, New Zealand, Oct 2023 - Sep 2024 and Oct 2025 - Sep 2025



Notes: Shows the difference of the matched pair between the start and end HoNOS65+ total score. Community admission to discharge includes discharge no further care and discharge other.

Interpretation: The graphs compare two time periods. Dark blue band indicates percentage improvement within the given time period, while black band indicates no significant change and light blue deterioration. Improvement = 4 or more, no significant change = -3 to 3 and deterioration = -4 or less.

Target: A greater percentage increase in improvement for both community and inpatient settings and a smaller percentage in deterioration.

Table 2: Difference in HoNOS65+ total score (12 items) for matched pairs by pair type and team, community New Zealand, Oct 2025 - Sep 2025


Team Type	Com Adm Rev				Com Rev Rev				Com Adm Dis			
	↓ Det	No SC	↑ Imp	N	↓ Det	No SC	↑ Imp	N	↓ Det	No SC	↑ Imp	N
Community team	13%	58%	29%	52	11%	78%	11%	650	11%	35%	54%	54
Kaupapa Māori team					8%	81%	11%	37				
Older people team	9%	62%	30%	466	11%	74%	15%	1,181	5%	51%	44%	928
Specialty team	8%	59%	34%	119	6%	88%	6%	294	2%	49%	49%	233
Total	9%	61%	30%	640	10%	78%	12%	2,197	5%	50%	46%	1,220

Notes: Shows the difference of the matched pair between the start and end HoNOS65+ total score. Community admission to discharge includes discharge no further care and discharge other.

Interpretation: Improvement (↑Imp) = 4 or more, no significant change (No SC) = -3 to 3 and deterioration (↓Det) = -4 or less.

Target: A greater percentage increase in improvement for community setting and a smaller percentage in deterioration.

Graph 3: Average number of clinically significant HoNOS65+ items by collection type: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025



Notes: Average number clinically significant items = the average number of items in the clinical range (2, 3 or 4) per collection. Community discharge does not include discharge to an inpatient unit.

Interpretation: Points are significantly different if error bars don't overlap. A decrease between admission and discharge is an indication of the outcomes achieved by the service and service user. A greater decrease between admission and discharge indicates a better outcome. A lower admission score could be indication of service users seeking out and being engaged by services at a lower level of severity.

Target: A greater decrease from admission to discharge and smaller percentage in clinical range at discharge


Table 3: Average number of clinically significant HoNOS65+ items by collection type and team: New Zealand, Oct 2025 - Sep 2025

Team type	Assessment only			Admission			Review			Discharge		
	N	Mean	CI	N	Mean	CI	N	Mean	CI	N	Mean	CI
Community services												
Alcohol and drug team							39	2.0	1.5-2.5			
Community team	349	3.6	3.4-3.8	201	3.8	3.5-4.1	1,708	2.7	2.7-2.8	166	2.3	2.0-2.6
Kaupapa Māori team							56	4.8	4.1-5.5			
Needs assessment and service coordination team							44	2.5	1.9-3.1			
Older people team	959	3.6	3.5-3.7	2,253	3.4	3.3-3.5	3,325	2.3	2.3-2.4	1,634	2.0	1.9-2.1
Pacific people team							26	1.6	1.0-2.2			
Specialty team	220	4.0	3.8-4.3	337	5.2	5.0-5.4	503	4.5	4.3-4.6	343	3.7	3.5-3.9
Total	1,541	3.6	3.5-3.7	2,816	3.6	3.5-3.7	5,714	2.7	2.6-2.7	2,160	2.3	2.2-2.4
Inpatient services												
Forensic team							25	2.3	1.3-3.2			
Inpatient team				336	4.4	4.1-4.7	44	3.1	2.2-4.0	219	2.5	2.2-2.8
Older people team				739	4.8	4.6-4.9	144	3.5	3.2-3.9	586	2.2	2.1-2.4
Total				1,084	4.6	4.5-4.8	213	3.3	3.0-3.6	814	2.3	2.2-2.5

Notes: N = number of collections in period. Average = average number of HoNOS65+ items in the clinically significant range (ie scoring 2, 3, or 4). CI = confidence interval for average score. Community discharge does not include discharge to an inpatient unit.

Interpretation: If confidence intervals for two scores do not overlap, then the scores have a statistically significant difference. It is important to note that statistical significance may not indicate a clinically significant difference.

Graph 4: Average number of clinically significant HoNOS65+ items at admission and discharge by ethnic group: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025




Notes: Average number of HoNOS65+ items in the clinical range (2, 3 or 4). Community discharge does not include discharge to an inpatient unit.

Interpretation: Points are significantly different if error bars don't overlap. Decrease between admission and discharge is an indication of the outcomes achieved by the service and service user. A greater decrease between admission and discharge indicates a better outcome. A lower admission score could be indication of service users seeking out and being engaged by services at a lower level of acuity.

Target: A greater decrease from admission to discharge and smaller percentage in clinical range at discharge.

Graph 5a: Percentage of collections in clinical range on each HoNOS65+ item (admission and discharge collections): New Zealand, Oct 2025 - Sep 2025



Notes: Percentage of service users in the clinical range (2, 3 or 4) for each HoNOS65+ item. Community discharge does not include discharge to an inpatient unit.

Interpretation: The longer the bar, the more prevalent the difficulty measured by the item is amongst service users. A greater decrease in the length of the bar from admission to discharge suggests a better outcome for the difficulty measured by the item. Items that show medium to high frequency and less change may suggest possible targets for service improvement.

Target: A greater decrease from admission to discharge and smaller percentage in clinical range at discharge.

Table 5a: Percentage of collections in clinical range on each HoNOS65+ item (admission and discharge collections) by team: New Zealand Oct 2025 - Sep 2025

Community services

First 6 HoNOS items														
Team type	N		AGR		SH		AOD		COG		PHY		DelHal	
	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch
Community team	201	166	20%	5%	15%	3%	9%	5%	41%	31%	56%	54%	30%	14%
Older people team	2,251	1,634	23%	7%	5%	0%	4%	2%	52%	49%	47%	43%	24%	11%
Specialty team	337	343	39%	21%	6%	2%	5%	3%	76%	79%	69%	66%	21%	17%
Total	2,813	2,160	25%	9%	6%	1%	5%	2%	54%	52%	51%	47%	24%	12%
Second 6 HoNOS items														
Team type	DEP		OTH		REL		ADL		LIV		OCC			
	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch
Community team	43%	14%	61%	33%	33%	16%	41%	31%	15%	4%	19%	17%		
Older people team	34%	7%	47%	20%	28%	14%	46%	40%	12%	3%	17%	6%		
Specialty team	49%	22%	67%	34%	49%	40%	66%	54%	29%	14%	44%	21%		
Total	37%	10%	51%	23%	30%	18%	48%	42%	14%	5%	21%	9%		

Inpatient services



First 6 HoNOS items														
Team type	N		AGR		SH		AOD		COG		PHY		DelHal	
	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch
Inpatient team	336	219	36%	11%	15%	3%	11%	6%	49%	42%	53%	38%	40%	24%
Older people team	738	586	47%	13%	19%	2%	10%	2%	58%	43%	48%	36%	50%	22%
Total	1,083	814	43%	12%	17%	3%	11%	4%	55%	43%	50%	36%	47%	22%
Second 6 HoNOS items														
Team type	DEP		OTH		REL		ADL		LIV		OCC			
	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch	Adm	Disch
Inpatient team	38%	17%	55%	32%	38%	22%	50%	27%	24%	13%	34%	16%		
Older people team	47%	19%	58%	26%	45%	17%	59%	31%	21%	6%	22%	8%		
Total	44%	18%	57%	28%	43%	18%	56%	30%	22%	8%	25%	11%		

Notes: Percentage of service users in the clinical range (2, 3 or 4) for each HoNOS65+ item. Community discharge does not include discharges to an inpatient unit.

Interpretation: The higher the percentage, the more prevalent the difficulty measured by the item is amongst service users. A greater decrease between admission and discharge suggests a better outcome for the difficulty measured by the item. Items that show medium to high frequency and less change may suggest possible targets for service improvement.

Target: A greater decrease from admission to discharge and smaller percentage in clinical range at discharge.

Graph 5b: Percentage of collections in clinical range on each HoNOS65+ item (review collections): New Zealand, Oct 2025 - Sep 2025



Notes: Percentage of service users in the clinical range (2, 3 or 4) for each HoNOS65+ item on review collections. Due to most admission being less than 91 days, the data set for review collections in inpatient settings is relatively small.

Interpretation: The longer the bar, the more prevalent the difficulty measured by the item is amongst service users.


Table 5b: Percentage of collections in clinical range on each HoNOS65+ item (review collections) by team: New Zealand, Oct 2025 - Sep 2025

Team type	N	AGR	SH	AOD	COG	PHY	DelHal	DEP	OTH	REL	ADL	LIV	OCC
Community services													
Alcohol and drug team	39	3%	0%	41%	10%	62%	13%	15%	31%	3%	5%	13%	5%
Community team	1,707	7%	3%	9%	30%	52%	27%	24%	42%	28%	30%	8%	17%
Kaupapa Māori team	56	20%	4%	11%	64%	80%	41%	41%	63%	48%	54%	27%	34%
Needs assessment and service coordination team	44	2%	2%	0%	34%	48%	16%	18%	27%	23%	39%	5%	39%
Older people team	3,325	8%	2%	3%	35%	40%	18%	25%	35%	20%	32%	6%	11%
Pacific people team	26	0%	0%	0%	19%	31%	27%	12%	8%	15%	15%	0%	32%
Specialty team	503	22%	1%	3%	70%	73%	17%	43%	57%	56%	56%	16%	32%
Total	5,713	9%	2%	5%	36%	47%	21%	26%	39%	25%	33%	8%	15%
Inpatient services													
Forensic team	25	32%	16%	0%	20%	32%	44%	4%	17%	24%	24%	4%	13%
Inpatient team	44	20%	5%	2%	27%	50%	30%	30%	45%	25%	36%	21%	23%
Older people team	144	30%	5%	8%	46%	48%	32%	38%	53%	28%	44%	11%	16%
Total	213	28%	6%	6%	39%	46%	33%	32%	47%	27%	40%	12%	17%

Notes: Percentage of service users in the clinical range (2, 3 or 4) for each HoNOS65+ item on review collections. Due to most admission being less than 91 days, the data set for review collections in inpatient settings is relatively small.

Interpretation: The higher the percentage, the more prevalent the difficulty measured by the item is amongst service users.

Graph 5c: Distribution of other mental health and behavioural problems in the clinical range by collection type: New Zealand, Oct 2025 - Sep 2025



Notes: This is for service users in the clinical range (2, 3 or 4) for other mental health and behavioural problems item.

Interpretation: This shows the distribution of what is selected for other mental health and behavioural problems. Other mental health and behavioural problems provides clinicians with the opportunity to rate the most severe clinical problem not covered elsewhere from the following list: phobic, anxiety, obsessive-compulsive, stress, dissociative, somatoform, eating, sleep, sexual, other. Other conforms to an ICD or DSM classification. The most severe clinical problem can change for each collection.


Table 5c: Distribution of other mental health and behavioural problems in the clinical range (admission and review collections) by team: New Zealand, Oct 2025 - Sep 2025

	Phobic	Anxiety	Obsessive-compulsive	Stress	Dissociative	Somatoform	Eating	Sleep	Sexual	Other	Number
Community service											
Community team	2%	46%	3%	21%	1%	1%	3%	18%	0%	5%	824
Kaupapa Māori team	0%	27%	12%	36%	0%	0%	3%	21%	0%	0%	33
Older people team	0%	52%	2%	9%	0%	1%	4%	23%	1%	8%	2,190
Specialty team	0%	61%	0%	13%	0%	1%	2%	16%	1%	5%	500
Total	1%	52%	2%	12%	0%	1%	3%	21%	1%	7%	3,582
Inpatient service											
Inpatient team	0%	42%	1%	16%	1%	0%	5%	25%	2%	7%	201
Older people team	0%	38%	2%	11%	1%	1%	8%	31%	2%	7%	493
Total	0%	39%	2%	12%	1%	1%	7%	29%	2%	7%	701

Notes: This is for admission and review collections. This is for service users in the clinical range (2, 3 or 4) for other mental health and behavioural problems item.

Interpretation: The shows the distribution of what is selected for other mental health and behavioural problems.

Graph 6: Index of Severity ratings for HoNOS65+ by collection type: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025




Notes: Community discharge does not include discharge to an inpatient unit. Index of Severity: Sub clinical = all items <2, mild = at least one item >1 and all items <3, moderate = at least one item >=3, severe = at least 2 items >=3 using first 10 items.

Interpretation: Darker bars indicate higher overall level of severity. More positive outcome is shown by larger decrease in darker sections of bar between admission and discharge.

2. Other measures of service activity

This section presents other information related to data collected in PRIMHD that may be helpful for understanding how teams are operating. This includes information relevant to caseload intensity and team activity.

Graph 7: Index of severity for HoNOS65+ (admission and review collections) by team, New Zealand, Oct 2025 - Sep 2025



Notes: Only data related to collection at admission and review is included so that results reflect the severity of service users during their engagement with the service. Index of Severity: Sub clinical = all items <2, mild = at least one item >1 and all items <3, moderate = at least one item >=3, severe = at least 2 items >=3 using first 10 items.

Interpretation: This graph gives an impression of the overall severity of the caseload for different teams at admission. The longer the darker bar, the higher the overall severity of the team’s caseload.


Table 7: Index of Severity for HoNOS65+ by collection type and team: New Zealand, Oct 2025 - Sep 2025

Team type	Assessment only					Admission					Review					Discharge				
	Sub	Mild	Mod	Sev	N	Sub	Mild	Mod	Sev	N	Sub	Mild	Mod	Sev	N	Sub	Mild	Mod	Sev	N
Community services																				
Alcohol and drug team											28%	38%	28%	5%	39					
Community team	3%	20%	30%	47%	349	5%	23%	29%	42%	201	14%	39%	24%	22%	1,708	25%	32%	20%	22%	166
Kaupapa Māori team											5%	23%	32%	39%	56					
Needs assessment and service coordination team											20%	41%	23%	16%	44					
Older people team	4%	25%	26%	45%	959	6%	25%	27%	42%	2,253	20%	34%	22%	24%	3,325	22%	27%	25%	26%	1,634
Pacific people team											35%	50%	12%	4%	26					
Specialty team	3%	17%	27%	53%	220	1%	12%	22%	66%	337	3%	26%	30%	41%	503	4%	27%	29%	40%	343
Total	4%	23%	27%	46%	1,541	5%	23%	27%	45%	2,816	17%	35%	24%	25%	5,714	20%	28%	25%	28%	2,160
Inpatient services																				
Forensic team											24%	28%	28%	20%	25					
Inpatient team						6%	22%	21%	50%	336	23%	32%	14%	32%	44	29%	30%	17%	24%	219
Older people team						3%	12%	18%	67%	739	12%	23%	19%	46%	144	26%	31%	21%	23%	586
Total						4%	15%	19%	62%	1,084	15%	25%	19%	40%	213	26%	31%	20%	23%	814

Notes: Sub = sub clinical, Mild = mild, Mod = moderate, Sev = severe on Index of Severity. Only admission collection data is included so that results reflect the severity of service users during their engagement with the service. Index of Severity: Sub clinical = all items <2, mild = at least one item >1 and all items <3, moderate = at least one item >=3, severe = at least 2 items >=3 using first 10 items. Community discharge does not include discharge to an inpatient unit.

Interpretation: Larger percentages in the columns to the right for each type of collection, the higher the overall severity of the team's caseload.

Graph 8: Collections with no HoNOS65+ items in clinical range: New Zealand, Oct 2025 - Sep 2025



Notes: Includes admission and review collections. Shows percentage of service users with all HoNOS65+ items less than two, ie no HoNOS65+ items in the clinical range.


Interpretation: There are a variety of reasons that may make it appropriate for service users to remain in the service even though they show no HoNOS65+ items in the clinical range. However, teams showing a larger or substantial percentage of service users with no HoNOS65+ items in the clinical range could benefit from reviewing these cases to ensure that the service remains appropriate for this service user.

Table 8: Collections with no HoNOS65+ items in clinical range: New Zealand, Oct 2025 - Sep 2025

Team type	Number of collections with no items in clinical range	Percentage with no items in clinical range
Community services		
Alcohol and drug team	11	22%
Community team	247	13%
Kaupapa Māori team	4	7%
Needs assessment and service coordination team	6	13%
Older people team	754	14%
Pacific people team	8	28%
Specialty team	16	2%
Total	1,049	12%
Inpatient services		
Forensic team	6	22%
Inpatient team	30	8%
Older people team	38	4%
Total	74	6%

Notes: For further information see notes for graph 8.

Graph 9: Focus of care categories: New Zealand, Oct 2023 - Sep 2024 and Oct 2025 - Sep 2025



Notes: Data from review and discharge collections.


Interpretation: Darker bars indicate more intensive involvement in care, so a longer, darker bar in general suggests more intensive working. A longer functional gain bar and shorter maintenance bar potentially suggests more recovery focused ways of working.

Table 9: Focus of care categories by team: New Zealand, Oct 2023 - Sep 2024 and Oct 2025 - Sep 2025

Team type	Acute		Functional gain		Intensive extended		Maintenance		Number of collections	
	Last	This	Last	This	Last	This	Last	This	Last	This
Community services										
Alcohol and drug team	0%	11%	12%	7%	7%	7%	81%	75%	58	55
Community team	6%	6%	12%	12%	3%	3%	79%	79%	1,794	1,665
Kaupapa Māori team	0%	2%	2%	8%	0%	4%	98%	86%	53	49
Needs assessment and service coordination team	2%	0%	8%	11%	2%	2%	89%	86%	66	44
Older people team	9%	9%	16%	15%	3%	3%	72%	73%	4,569	4,889
Pacific people team	0%	4%	48%	15%	0%	0%	52%	81%	23	26
Specialty team	11%	15%	16%	15%	9%	13%	63%	57%	757	743
Total	8%	9%	15%	14%	4%	4%	73%	73%	7,346	7,492
Inpatient services										
Forensic team		7%		0%		7%		86%		29
Inpatient team	69%	69%	5%	11%	3%	2%	23%	18%	210	245
Older people team	72%	71%	0%	3%	1%	1%	27%	26%	651	691
Total	70%	69%	1%	5%	2%	1%	27%	25%	878	970

Notes: For further information see notes for Graph 9.

Graph 10: HoNOS65+ total score (review collections) by focus of care: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025



Notes: This data is just for review collections.

Interpretation: Points are significantly different if error bars don't overlap. A general downward trend in scores from acute to maintenance focus of care might be expected.

Table 10: HoNOS65+ total score (review collections) by focus of care by team: New Zealand, Oct 2025 - Sep 2025


Team type	Acute			Functional Gain			Intensive Extended			Maintenance		
	N	Mean	CI	N	Mean	CI	N	Mean	CI	N	Mean	CI
Community services												
Alcohol and drug team										32	6.3	4.8-7.8
Community team	50	14.4	12.7-16.1	162	8.5	7.8-9.2	45	12.8	10.9-14.6	1,216	8.7	8.4-9.0
Kaupapa Māori team										41	17.0	15.3-18.6
Needs assessment and service coordination team										37	8.3	6.7-9.8
Older people team	114	9.4	8.4-10.4	460	7.6	7.2-8.0	118	10.2	9.1-11.3	2,516	7.6	7.4-7.8
Pacific people team										20	5.5	3.6-7.3
Specialty team	41	14.5	13.0-16.0	86	12.3	11.3-13.3	72	13.0	11.9-14.2	287	13.9	13.3-14.4
Total	208	11.6	10.8-12.4	725	8.3	8.0-8.7	241	11.5	10.8-12.3	4,159	8.4	8.2-8.6
Inpatient services												
Forensic team										22	6.9	4.9-8.9
Older people team	97	13.5	12.2-14.8							37	9.2	7.1-11.3
Total	112	13.0	11.8-14.2							78	8.8	7.4-10.2

Notes: For further information see notes for Graph 10.

3. Collection completion and validity

This section presents information about the completeness and validity of the data on which the remainder of this report is based. It also shows the current targets for the variables presented.


Graph 11: Percentage of service users with at least one collection during the period in a quarter: New Zealand, Oct 2024 – Sep 2025 (65 years and over)



Interpretation: The longer the dark part of the bar, the more completely the data set includes all service users, and the more meaningful and representative the graphs, tables, and analyses using these data will be. This data is approximate due to movement of service users between teams and similar variations; however, it does provide a reasonably accurate representation of completion of measures. Only data with valid collections is included.


Target: To meet or exceed the target shown on the graph for the percentage of service users with at least one collection within the period.

Graph 12: Percentage of service users with admission and discharge collections completed by quarter: New Zealand, Oct 2024 – Sep 2025 (65 years and over)



Interpretation: The longer the dark bar, the larger the percentage of admission and discharge that had the relevant data collection type. Data is approximate due to movement of service users between teams and similar variations; however, it does provide a reasonably accurate representation of completion of measures. Only data with valid collections is included.

Graph 13: Percentage of valid collections, HoNOS65+: New Zealand, Oct 2023 - Sep 2024 and Oct 2024 - Sep 2025



Notes: Valid = Two or fewer of the 12 items scored as unknown or missing. Discharge excludes collection types for lost to care, discharge dead and brief episode of care. Collection in drug and alcohol teams is not required, therefore AOD data is excluded.

Interpretation: The longer the dark lines, the higher the percentage of valid scores.

Target: Aim for 95% valid collections.

Table 13: Invalid collections by team, by HoNOS65+ item: New Zealand, Oct 2025 - Sep 2025

Team type	% Invalid			Number of invalid ratings by HoNOS item												Total number
	Adm	Rev	Dch	1	2	3	4	5	6	7	8	9	10	11	12	
Community services																
Community team	2%	1%	12%	48	53	63	61	50	55	54	58	56	58	61	100	2,558
Kaupapa Māori team		0%		0	0	0	0	0	0	0	2	0	0	0	0	62
Needs assessment and service coordination team		0%		0	0	0	0	0	0	0	0	0	0	0	0	59
Older people team	1%	1%	9%	208	216	230	271	220	320	298	311	241	240	265	289	8,697
Pacific people team		0%		0	0	0	0	0	0	0	0	0	0	0	2	31
Specialty team	0%	1%	2%	8	10	11	10	7	25	17	16	14	11	22	20	1,436
Total	1%	1%	8%	265	280	305	343	278	401	370	388	312	310	351	414	12,872
Inpatient services																
Forensic team		0%		0	1	2	1	1	1	1	2	1	1	3	2	32
Inpatient team	4%	2%	2%	7	3	12	9	10	15	16	16	22	9	22	32	615
Older people team	5%	3%	4%	15	25	37	27	22	33	30	45	54	28	165	176	1,529
Total	4%	3%	3%	22	29	51	37	33	49	47	63	77	38	190	210	2,176

Notes: Per cent (%) invalid collections = the percentage of collections that had three or more of the 12 items scored as unknown or missing. Number of invalid ratings by HoNOS65+ item = for each of the HoNOS65+ items, the number of collections for which that item was unknown or missing. Discharge excludes collection types for lost to care, deceased and brief episode of care. Collection in drug and alcohol teams is not required, therefore AOD data is excluded.

Interpretation: The lower the per cent invalid, the higher the percentage of valid scores. The lower the number of invalid collections by HoNOS65+ item, the more collections that have valid data on that HoNOS65+ item.

Target: Aim for 95% valid collections (5% invalid collections) or better. Aim for as few invalid items as possible, with all items having a similar validity rate.